

STAGED SURGICAL APPROACH TO THE TREATMENT OF PATIENTS WITH THYROID CYSTS

Timur B. Makhmudov ¹, Ulugbek A. Sherbekov ²

1 PhD, Assistant of the Department of General Surgery,
Samarkand State Medical University, Samarkand, Uzbekistan
E-mail: dr.timur.maxmudov@gmail.com

2 DSc, Head of the Department of General Surgery,
Samarkand State Medical University, Samarkand, Uzbekistan
E-mail: ulasher68@gmail.com

ABSTRACT

We analyzed the surgical treatment of 128 patients with thyroid cysts who underwent surgery due to the presence of cysts measuring 3 centimeters or more in diameter. The surgical treatment of thyroid cysts was analyzed, indications and contraindications for this treatment method were determined, and their effectiveness was assessed. The developed algorithm allows you to choose a treatment method and management tactics for patients with thyroid cysts, depending on the size of the cyst and its morphological picture.

Key words: thyroid cyst, management tactics, surgical treatment.

INTRODUCTION

Relevance of the topic: Thyroid cyst (TC) refers to nodular formations of the thyroid gland with a cavity that is filled with liquid contents, determined by palpation or using imaging studies (1, 3). Thyroid cysts (nodes) can be asymptomatic for a long time, and when they increase in size, they cause “compression syndrome” of nearby neck structures, disrupting the functioning of the thyroid gland, leading to suffocation, hoarseness, difficulty swallowing and painful sensations (2, 5). A thyroid cyst is a small but common benign tumor with a cavity containing fluid. Such a lesion occurs in almost 50% of the planet's inhabitants, and in people of pre-retirement and retirement age, this pathology is considered practically the norm (4, 6, 7). The most dangerous complications of cysts are inflammation and suppuration of the nodes, malignant degeneration (8,

11). Nodules of various morphological forms can develop in the thyroid gland, and most of them are benign in nature (nodular colloid goiter, adenoma, thyroid cyst (9, 10). Cysts (nodules) are the most common pathology of the thyroid gland, and in the female population it occurs 4-8 times more often (12,13). Despite the widespread prevalence of this disease and various approaches to the treatment of this pathology, the issue of surgical tactics for managing patients with thyroid cysts, depending on their size and morphological forms, remains relevant.

Purpose of the study: Conduct a factor analysis and develop an approach to surgical treatment of thyroid cysts depending on the morphological structure and size of the cyst.

Material and research methods: From 2015 to 2023, 128 patients with thyroid cysts underwent surgical treatment in the surgical departments of the clinic of SamSMU and SamSMO.

The age of the patients ranged from 16 to 78 years, most of the patients were of working age from 41 to 60 years. The distribution of patients by gender and age in the study groups was identical.

The degree of enlargement of the thyroid gland was assessed according to the classification of Nikolaev O.V. (Figure 1), based on ultrasound and palpation of the thyroid gland. In 83 (64.8%) patients, the examination revealed III degree enlargement of the thyroid gland, 39 (30.5%) patients - IV and 6 (4.7%) - V degree. Depending on the location of the cyst, it was detected in the left or right lobe in 63 (49.3%) patients, in both lobes in 48 (37.5%) and in the isthmus in 17 (13.2%) (Table 1).



Figure. 1. Patient M., 68 years old. Stage V thyroid cyst.

Table 1**Distribution of patients depending on the degree of enlargement of the thyroid gland and the location of the cysts.**

Options	III	IV	V	Total
The degree of enlargement of the t/g	64,8%	30,5%	4,7%	100%
Localization of cysts	In one of the lobes	In both lobes	isthmus	
	49,3%	37,5%	13,2%	100%

All patients underwent a comprehensive examination, which included laboratory tests (general blood count, urine test, biochemical blood test, prothrombin index and time, blood ELISA), the level of thyroid hormones (T3, T4 and TSH, antibodies to thyroglobulin and thyroid peroxidase) was determined at baseline and at various times after treatment. Instrumental diagnostic methods were used (chest x-ray, ultrasound (Figure 2), MRI (Figure 3).




Figure 2. Patient K., 51 years old. On the echogram, there is a cyst of the right lobe of the thyroid gland, degree II

When conducting instrumental studies, we paid great attention to the nature of the cavity formation; simple cavity formations containing a liquid component or a colloid or a solid (tissue/epithelial) component were interpreted in most cases as a benign formation that has the lowest risk of degeneration. Complex formations, multi-chambered in nature with a solid component, were regarded as formations with a high possibility of malignant degeneration.




Figure 3. Patient V., 68 years old. MRI shows a grade V thyroid cyst, with compression of the neck organs.

All patients undergoing surgical treatment underwent morphological examination as in the preoperative period, which included fine-needle aspiration biopsy (FNA), intraoperative express biopsy and planned postoperative histological examination of removed thyroid tissue. In the postoperative period, a comparison was made (in 128 patients) of the FNA data and the final histological examination of the removed thyroid tissue in order to determine the likelihood of malignancy of the node. Based on the data obtained, the informative value of FNA at the preoperative stage was 89.9%.

We studied the results of histological reports of all operated patients (n=128). The following histological forms of thyroid cysts were determined: - colloid cyst in 75 (58.6%) patients; - follicular cyst in 46 (35.9%) patients; - cystadenom in 7 (5.5%) patients (Table 2).

Table 2
Morphological nature of thyroid cysts in a group of patients undergoing surgical treatment.

Morphological picture	Number of patients	
	Num	%
Colloid cyst	75	58,6
Follicular cyst	46	35,9
Cystadenom	7	5,5
Total	128	100,0

Patients (n=128) with cyst sizes of 3 cm or more underwent surgical removal of cysts (Figure 4). In the presence of a unilateral lesion and a positive

morphological picture, we performed hemistrulectomy in 62 patients (48.4%). In the presence of cystic lesions of both lobes and a positive morphological picture, subtotal thyroidectomy was performed, which was performed in 52 (40.6%) patients; and in the case of a negative morphological picture, regardless of the number of cysts and their location, total thyroidectomy was performed in 14 (11%) patients (Table 3).




Figure 4. Patient Sh., 56 years old. Hemorrhage into the cyst cavity after minimally invasive treatment. An increase in the size of the cyst with symptoms of compression of the neck organs.

Table 3

The number and nature of surgical interventions performed, depending on the location of the cysts and the morphological picture.

Location of cysts and morphological picture	Nature of surgical interventions			
	Hemistrulectomy	Subtotal strumectomy	Total strumectomy	%
In one of the lobes with a positive morphological picture	62	-	-	48,4%
In one of the shares with a negative morphological picture	-	-	5	4%
In both lobes with positive morphological picture	-	52	-	40,6%
In both lobes with a negative morphological picture	-	-	9	7%
Total	128			100%

In the long-term postoperative period, relapse of the disease was observed in the group of patients who underwent hemistrumectomy in 2 cases, which amounted to 3.2% of the total number of patients who underwent this operation. After subtotal and total strumectomy, no relapses were observed.

Results and discussion. In a group of patients who underwent surgical treatment in the period from 2015 to 2023, an analysis of the ultrasound and MRI structure of the cavity structure of the thyroid gland was carried out. Of the 128 patients in this group, 86 were diagnosed with simple cysts, and 42 with complex, multilocular formations. Analysis of histological examination of complex cavity formations showed the presence of 33.3% of cases of their malignancy.

Comparative analysis of preoperative FNA with data from postoperative histological studies. The effectiveness of TAPB in patients with thyroid cysts was 89.9%.

Analysis of the morphological structure showed a predominance of colloid forms of cysts (58.6%), over follicular (35.5%) and cystadenomas (5.5%), respectively.

In the group of patients who underwent surgical treatment, the largest number of patients underwent hemistrumectomy (48.4%), the second most performed was subtotal strumectomy (40.6%), total strumectomy accounted for 11% of the total number of surgical interventions.


Relapse of the disease was observed only in the group of patients who underwent hemistrumectomy and amounted to 3.2%.

Based on our research, we have developed an algorithm for the management of patients with thyroid cysts (Table 4):

Conclusions.

1. Surgical removal of thyroid cysts is the most effective and at the same time radical treatment method used in the development of cyst complications.
2. Staged diagnostic and morphological studies of thyroid cysts allow you to choose the tactics of surgical intervention.
3. Instrumental diagnosis of the structure of cysts allows early differentiation between simple cavitory formations and complex multi-chamber formations and choosing the optimal method of treating cysts.
4. The developed diagnostic and treatment algorithm for choosing a treatment method for thyroid cysts, based on stage-by-stage diagnostic and morphological studies, has improved the results of treatment of this disease.

Table 4

Algorithm for the management of patients with thyroid cysts

REFERENCES

1. Belyakov I.E. and others. Comprehensive assessment of nodular pathology of the thyroid gland // Health and education in the XXI century. – 2017. – T. 19. – No. 8.
2. Beltsevich D.G., Vanushko V.E., Melnichenko G.A., Rumyantsev P.O., Fadeev V.V. Clinical recommendations of the Russian Association of Endocrinologists for the diagnosis and treatment of multinodular goiter in adults. Project. — 2016.
3. Braverman L.I., Diseases of the thyroid gland. Moscow, Medicine, 2000.
4. Carle, A., Andersen, S. L., Boelaert, K. & Laurberg P. Management of endocrine disease: subclinical thyrotoxicosis: prevalence, causes and choice of therapy. Eur. J. Endocrinol. 176, R325–R337 (2017).
5. Castellnou S, Lifante JC, Polazzi S, Pascal L, Borson-Chazot F, Duclos A. Influence of Care Pathway on Thyroid Nodule Surgery Relevance: A Historical Cohort Study. J Clin Med. 2020 Jul 17;9(7):2271.
6. Elsayed AA, Murdoch C, Murray S, et al. Can thyroid surgery be decided based on ultrasonographic findings, irrespective of cytopathological findings? Five-year retrospective study in a district general hospital. Clin Radiol 2017;72:170–4.
7. Jackson BS. Controversy regarding when clinically suspicious thyroid nodules should be subjected to surgery: Review of current guidelines. Medicine (Baltimore). 2018 Dec;97(50):e13634.
8. Reinisch A, Malkomes P, Habbe N, et al. Guideline compliance in surgery for thyroid nodules—a retrospective study. Exp Clin Endocrinol Diabetes. 2017;125:327–334.
9. Sleptsov I.V. Thyroid nodules. Modern principles of diagnosis and treatment. - M.: Publishing house "Elite", 2014. - 96 p.
10. Ushakov A.V. Benign diseases of the thyroid gland. Clinical classification. - M.: Clinic of Dr. A.V. Ushakova, 2013. - 384 p.
11. Vanushko, V.E. Nodular goiter (clinical lecture) / V.E. Vanushko, V.V. Fadeev // Endocrine surgery. – 2012. – No. 3. – P. 11–16.
12. Wolinski, K. Usefulness of different ultrasound features of malignancy in predicting the type of thyroid lesions: a meta-analysis of prospective studies / K. Wolinski, M. Szkudlarek, E. Szczepanek-Parulska, M. Ruchala // Pol. Arch. Med. Wewn. – 2014. – Vol. 124, № 3. – P. 97–104.
13. Zefirova G.S., Diseases of the thyroid gland., Moscow, 1999.